



Namibia onshore Hydrocarbon Prospectivity

Ansgar Wanke University of Namibia Petroleum Geology

Purpose of Talk

Oshana

Omusat

Cunene



North West



South

1. Overview of Namibian Onshore Basins

Kavango

2. New encouraging insights

Oshikoto

3. How and why to Explore



Cubango

Namibia 825 419 km²

Two groups of onshore basins:

- Neoproterozoic-Early Cambrian riftforeland basins
- Karoo (rift) basins (Upper Carboniferous – Jurassic)

Onshore basins cover some 350 000km² > 40% of the country

12 onshore wells with TD between 600-2500m



Neoproterozoic / Early Cambrian Basins

- 1: Owambo-Etosha Basin
- 2: Nama Basin
- Basins are bordered by early Cambrian Orogenic Belts
- Rift, Platform and Molasse deposits
- Foreland Basin architecture
- Partly covered by Karoo succession
- Largely covered by a veneer of Cainozoic sediments





Owambo Basin

Vintage Exploration

- Seismic campaigns 1969-1995
- 12 wells of which 5 HC exploration wells (1964-1986)

Exploration from 2000

- Four aerial surveys (mag/gravity) (2004, 11, 13, 14)
- Soil gas sampling (2012)
- 120km 2D test lines (2017)
- Airbus remote sensing study (2018)
- Passive seismic (2018)

Findings

- Delineation of depocenters mature fairways
- Anticlinal and roll over traps towards basin margin
- · Carbonate built-ups as potential reservoirs
- Intriguing correlation of structures, soil gas HCs, and passive seismic for DHSs

Well	Year	Operator	TD (m)	Well Result
ST-1	1964	Texas Eastern	1875	Dry Hole
Etosha 1-1	1970	Brilund	1584	Dry Hole
Etosha 2-1	1970	Brilund	1228	Dry Hole
Etosha 5-1A	1970	Brilund	2509	Oil shows
OPO-1	1986	OPIC	700	Dry Hole



Summary of exploration history in the Owambo Basin until 2017. Modified from Hoak et al. (2014)



1986)

Owambo Basin Geology

Tectonostratigraphy

- Syn-rift followed by passive margin platform deposition, followed by retroarc foreland basin formation with molasse deposition during Damara (pan-African) orogeny
- Neoproterozoic to early Cambrian deposition
- **Compression during Cambrian** with minor Phanerozoic reactivation events

Wells

- Three wells pass through Black Shale marker (2.8% TOC)
- Only one exploration well reaches top of platform sequence at TD 2509m
 - Two deeper source rock levels expected, better chance of maturity
- Sandstones of good RQ and fractured carbonates intersected (<15% porosity)
- HC show in one well



Schematic structural S-N cross-section through the Owambo Basin. Modified from Hoak et al. (2014)

MODIFIED AFTER YPMA, 1978

Recent Findings

Structural Modelling

- Structures modelled with magnetic-gravity near basin margin
- Reveals foldbelt with larger • anticlines
- Consistent with seismic



op Tsumeb Carbo

Gravimetric structural interpretation coincides largely with magnetic interpretation

Widely spaced 2D seismic showing shallow top platform carbonates (red) coinciding with anticlines deduced from gravitymagnetics



Aeromagnetic structural interpretation of the southern Owambo Basin. High frequency have been filtered using upward continuation from 2000m, reduced to pole.



Consulting NAMIBIA

Nama Basin

- Geologically roughly the southern counterpart to the Owambo Basin
 - Less knowledge on deeper parts of the basin
- Long exploration history inspired by surface oil shows
- 5 wells drilled (1928, 63, 68, 92), all
 <2300m TD
- Aeromagnetic (1992)
- 360 km vintage 2D seismic (1968)
- 500 km 2D (2008)
 - Indicates an at least 7km deep basin containing three megasequences
- Geochemical studies on oil shows (Summons et al., 2008)





Map of main foreland basins, adapted from Bray & Lawrence (1998)

Karoo Basins

Depocenters defined by:

- 1. Sag/rift Basins
- 2. Over-deepened glacial valleys
- 3. Transfer basins
- Most basins shallow
 - < 1000m,
 - CBM potential, no or limited conventional potential
- Transfer Basins potentially deep
 - High conventional and unconventional potential
 - Lokichar style analogue?



Transfer and Rift Basins in Southern Africa and Namibia

MARIMBA sediment thickness inversion



Southern African Tectonic Elements SW-NE cross section underlain by MARIMBA sediment thickness inversion (Granath & Dickson, 2018)









Cross section through the Lokichar Basin, Kenia (Morley at al. 1999).

Conclusions



- Namibia > 40% covered by basins
- Two large Neoproterozoic-Cambrian Basins with HC indicators
 - Geophysics indicate deep depocenters and structural/stratigraphic closures
- Deep transfer basins with Karoo fill likely hidden under Cainozoic cover
 - Localized pull-apart/releasing bend basins hosting thick Permian Jurassic packages
 - · Could be similarly prospective as East African rift basins of similar size
- How to explore?
 - High resolution gravimetry/magnetics
 - Low cost HC indicators (soils gas, passive seismic, etc.)
 - Stratigraphic wells > 3000m, thereafter seismic
- Why?
 - Relatively low costs to start
 - Strong Local geoscience expertise support
 - Easy field access









Pioneer Oil & Gas Consulting NAMIBIA

G & G on-/offshore License Applications Local Content

yuri.perez@pioneerenergy.com.na

THANK YOU